DESTINED SENTER

JUL 2 8 2006

Page 3

Application No.: 10/759898 Docket No.: CL2035USNA

Amendments to Claims

Claim 1 (Withdrawn). A method of increasing the yield of an aromatic carboxylic acid from a host cell producing said aromatic carboxylic acid comprising:

- a) providing a host cell which:
 - i) produces an aromatic carboxylic acid; and
- ii) comprises at least one yhcQ gene and at least one yhcP gene; and
 b) up-regulating the expression of the at least one yhcQ gene and the at least one yhcP gene whereby the yield of aromatic carboxylic acid is increased.

Claim 2 (Currently Amended). A method for increasing the resistance of a host cell to aromatic carboxylic acids comprising:

- a) providing a host cell which comprises at least one <u>E. coli_yhcQ</u> gene and at least one <u>E. coli_yhcP</u> gene; and
- b) up-regulating the expression of the at least one <u>E. coli yhcQ</u> gene and the at least one <u>E. coli yhcP</u> gene whereby the host cell resistance to aromatic carboxylic acids is increased.
- Claim 3 (Currently Amended). A method according to either Claim 1 or Claim 2 wherein the at least one yhcQ gene and the at least one yhcP gene are endogenous to said host cell.
- Claim 4 (Currently Amended). A method according to either Claim 1 or Claim 2 wherein the at least one yhcQ gene and the at least one yhcP gene are heterologous to said host cell.
- Claim 5 (Currently Amended). A method according to either Claim 1 or Claim 2 wherein the host cell is selected from the group consisting of bacteria, yeast, fungi and plants.
- Claim 6 (Original). A method according to Claim 5 wherein the host cell is an enteric bacteria.
- Claim 7 (Original). A method according to claim 5 wherein the host cell is selected from the group of genera consisting of Escherichia, Salmonella, Bacillus, Acinetobacter, Streptomyces, Methylobacter, Rhodococcus, Corynebacterium, Pseudomonas, Rhodobacter, and Synechocystis.
- Claim 8 (Currently Amended). A method according to either Claim 1 or Claim 2 wherein the aromatic carboxylic acid is selected from the group consisting of of para-

Application No.: 10/759898 Docket No.: CL2035USNA

JUL. 28. 2006 12:37PM

Page 4

hydroxybenzoic acid, para-hydroxycinnamic acid, cinnamic acid, salicylic acid, benzoic acid, and 1-napthoic acid.

Claim 9 (Currently Amended). A method according to either Claim 1 or Claim 2 wherein the at least one yhcQ gene has the nucleic acid sequence as set forth in SEQ IDNO:2 and the at least one yhcP gene has the nucleic acid sequence as set forth in SEQ IDNO:1. comprise at least one suitable promoter operably linked to at least one isolated nucleic acid molecule selected from the group consisting of SEQ ID NO:1—4.

Claim 10 (Canceled).

Claim 11 (Currently Amended). A method according to either Claim 1 wherein the at least one yhcQ gene and the at least one yhcP gene are expressed on a multicopy plasmid.

Claim 12 (Currently Amended). A method according to either Claim 1 or Claim 2 wherein the at least one yhcQ gene and the at least one yhcP gene are under the control of a strong promoter selected from the group consisting of lac, trp, lP_L, lP_R, T7, tac, and trc.

Claim 13 Canceled).

- Claim 14 (Withdrawn). A chimeric gene encoding an efflux protein comprising:
 a) an isolated nucleic acid molecule having a nucleic acid sequence selected from the group consisting of:
 - i) an isolated nucleic acid molecule selected from the group consisting of SEQ ID NO:1-4; and
 - ii) an isolated nucleic acid molecule, wherein said isolated nucleic acid molecule hybridizes with (i) under the following conditions: 0.1×SSC, 0.1% SDS at 65°C and washed with 2×SSC, 0.1% SDS followed by a second wash in 0.2×SSC, 0.1% SDS; and
- b) a promoter;

wherein the promoter is heterologous to the isolated nucleic acid molecule.